


Combining interval analysis and nonlinear predictive control to compute capture tubes


L. Jaulin, J. Ninin, G. Chabert,
S. Le Menec, M. Saad,
V. Le Doze, A. Stancu
April, 3, 2014, Paris, CNAM.


1 V-stability



Vaimos (IFREMER and ENSTA)








X_1 : outside the corridor.


X_2 : inside the corridor.

Definition. Consider a differentiable function $V(\mathbf{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$. The system is V -stable if

$$\left(V(\mathbf{x}) \geq 0 \Rightarrow \dot{V}(\mathbf{x}) < 0 \right).$$

Checking the V -stability can be done using interval analysis.





Système non-holonomes


2 Station keeping problem

The problem of *station keeping* for a robot is to stay inside a disk around origin.

Consider a non holonomous robot described by

$$\begin{cases} \dot{x} &= \cos \theta \\ \dot{y} &= \sin \theta \\ \dot{\theta} &= u. \end{cases}$$

Since $\dot{x}^2 + \dot{y}^2 = 1$, this robot cannot stop.



Transformation from Cartesian to polar

The polar form for the state equations is


$$\left\{ \begin{array}{ll} \text{(i)} & \dot{\varphi} = \frac{\sin \varphi}{d} + u \\ \text{(ii)} & \dot{d} = -\cos \varphi. \\ \text{(iii)} & \dot{\alpha} = -\frac{\sin \varphi}{d}. \end{array} \right.$$


We propose here the following control




$$u = \begin{cases} +1 & \text{if } \cos \varphi \leq \frac{1}{\sqrt{2}} \quad (\text{the robot turns left}) \\ -\sin \varphi & \text{otherwise} \quad (\text{the robot goes toward zero}) \end{cases}$$



The closed loop state equations are

$$\left\{ \begin{array}{l} \text{(i)} \quad \dot{\varphi} = \begin{cases} \frac{\sin \varphi}{d} + 1 & \text{if } \cos \varphi \leq \frac{1}{\sqrt{2}} \\ \left(\frac{1}{d} - 1\right) \sin \varphi & \text{otherwise} \end{cases} \\ \text{(ii)} \quad \dot{d} = -\cos \varphi. \end{array} \right.$$









3 Capture tubes

Consider the time dependant system


$$\mathcal{S} : \dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, t)$$

and a *tube*

$$\mathbb{G}(t) \subset \mathbb{R}^n, t \in \mathbb{R}.$$

The tube $\mathbb{G}(t)$ is said to be a *capture tube* if

$$\mathbf{x}(t) \in \mathbb{G}(t), \tau > 0 \Rightarrow \mathbf{x}(t + \tau) \in \mathbb{G}(t + \tau).$$



Theorem. Consider the tube

$$\mathbb{G}(t) = \{\mathbf{x}, \mathbf{g}(\mathbf{x}, t) \leq \mathbf{0}\}$$


where $\mathbf{g} : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^m$. If the system

$$\left\{ \begin{array}{ll} \text{(i)} & \underbrace{\frac{\partial g_i}{\partial \mathbf{x}}(\mathbf{x}, t) \cdot \mathbf{f}(\mathbf{x}, t) + \frac{\partial g_i}{\partial t}(\mathbf{x}, t)}_{\dot{g}_i(\mathbf{x}, t)} \geq 0 \\ \text{(ii)} & g_i(\mathbf{x}, t) = 0 \\ \text{(iii)} & \mathbf{g}(\mathbf{x}, t) \leq \mathbf{0} \end{array} \right.$$

is inconsistent for all (\mathbf{x}, t, i) , then $\mathbb{G}(t)$ is a capture tube for $\mathcal{S} : \dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, t)$

A software Bubbibex has been built by students from ENSTA Bretagne.

Bubbibex uses interval analysis to prove the inconsistency.



4 Test-case

Robot

$$\begin{cases} \dot{x} &= u_1 \\ \dot{y} &= u_2 \\ \dot{\theta} &= -\theta. \end{cases}$$

Target $(x_d, y_d) = (t, 0)$. We choose the control

$$u_1 = -x + t, \quad u_2 = -y.$$

Remark. We have

$$\dot{x} = -x + t$$

i.e.

$$x(t) = e^{-t}(x_0 + 1) + t - 1$$

The error on x is

$$e_x(t) = e^{-t}(x_0 + 1) - 1$$

The closed loop system satisfies

$$\begin{cases} \dot{x} &= -x + t \\ \dot{y} &= -y \\ \dot{\theta} &= -\theta. \end{cases}$$

Target tube. The tube we want is

$$\mathbb{G}(t) = \{\mathbf{x} \mid \mathbf{g}(\mathbf{x}, t) \leq \mathbf{0}\},$$

with

$$\begin{cases} g_1(\mathbf{x}, t) &= (x_1 - t)^2 + x_2^2 - r^2 \\ g_2(\mathbf{x}, t) &= (\cos x_3 - 1)^2 + \sin^2 x_3 - 0.2. \end{cases}$$

For $r = 4$, Bubbibex proves that $\mathbb{G}(t)$ is a capture tube.

For $r < 1$, some trajectories leave $\mathbb{G}(t)$ forever.

5 Lattice and capture tubes

Consider $\mathcal{S} : \dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, t)$.

If \mathbb{T} is the set of tubes and \mathbb{T}_c is the set of all capture tubes of \mathcal{S} then (\mathbb{T}_c, \subset) is a sublattice of (\mathbb{T}, \subset) .

We have indeed

$$\left\{ \begin{array}{l} \mathbb{G}_1(t) \in \mathbb{T}_c \\ \mathbb{G}_2(t) \in \mathbb{T}_c \end{array} \right. \Rightarrow \left\{ \begin{array}{l} \mathbb{G}_1(t) \cap \mathbb{G}_2(t) \in \mathbb{T}_c \\ \mathbb{G}_1(t) \cup \mathbb{G}_2(t) \in \mathbb{T}_c \end{array} \right.$$

Remark. If $\mathbb{G}(t) \in \mathbb{T}$, define

$$\text{capt}(\mathbb{G}(t)) = \bigcap \left\{ \overline{\mathbb{G}}(t) \in \mathbb{T}_c \mid \mathbb{G}(t) \subset \overline{\mathbb{G}}(t) \right\}.$$

This set corresponds to the smallest capture tube which encloses $\mathbb{G}(t)$.

6 Computing capture tubes

Problem. Given $\mathbb{G}(t) \in \mathbb{T}$, compute an interval $[\mathbb{G}^-(t), \mathbb{G}^+(t)] \in \mathbb{IT}$ such that

$$\text{capt}(\mathbb{G}(t)) \in [\mathbb{G}^-(t), \mathbb{G}^+(t)] .$$

Proposition 1. We have


$$\text{capt}(\mathbb{G}(t)) = \{(\mathbf{x}, t) \mid \exists (\mathbf{x}_0, t_0), \mathbf{x}_0 \in \mathbb{G}(t_0) \\ t \geq t_0, \mathbf{x} = \phi_{t-t_0}(\mathbf{x}_0, t_0) \}.$$


Proposition 2. We have

$$\text{capt}(\mathbb{G}(t)) = \mathbb{G}(t) \cup \Delta\mathbb{G}(t),$$

with

$$\Delta\mathbb{G}(t) = \left\{ (\mathbf{x}, t) \mid \begin{array}{l} \exists (\mathbf{x}_0, t_0), \mathbf{x}_0 \in \partial\mathbb{G}(t_0), \\ t \geq t_0, \mathbf{x} = \phi_{t-t_0}(\mathbf{x}_0, t_0) \\ \phi_{]t_0, t]}(\mathbf{x}_0, t_0) \notin \mathbb{G}(t) \end{array} \right\}$$






7 Test case


$$\mathcal{S} : \begin{cases} \dot{x} &= -x + t \\ \dot{y} &= -y \\ \dot{\theta} &= -\theta. \end{cases}$$


$$\mathbb{G}(t) : \begin{cases} g_1(\mathbf{x}, t) &= (x_1 - t)^2 + x_2^2 - r(t) \\ g_2(\mathbf{x}, t) &= (\cos x_3 - 1)^2 + \sin^2 x_3 - 0.2. \end{cases}$$


$$r(t) = 0.2 \cdot (t + 1)^2.$$


Some trajectories leave and come back to $\mathbb{G}(t)$.











References

F. Lydoire, P. Poignet. Nonlinear Model Predictive Control via Interval Analysis, IEEE Conference on Decision and Control, 2005. CDC'05.

L. Jaulin and F. Le Bars (2012). An interval approach for stability analysis; Application to sailboat robotics. IEEE Transaction on Robotics.

L. Jaulin(2014). Outer approximation of attractors using an interval quantization. Reliable Computing, Accepted,