

Interval robotics

Chapter 6: SLAM


Luc Jaulin,

ENSTA-Bretagne, Brest, France

<http://www.ensta-bretagne.fr/jaulin/>

1 Basic SLAM

$$\begin{cases} \dot{\mathbf{x}} &= \mathbf{f}(\mathbf{x}, \mathbf{u}) & \text{(evolution equation)} \\ \mathbf{y} &= \mathbf{g}(\mathbf{x}, \mathbf{u}) & \text{(observation equation)} \\ \mathbf{z}_i &= \mathbf{h}(\mathbf{x}, \mathbf{u}, \mathbf{m}_i) & \text{(mark equation)} \end{cases}$$



Redermor, GESMA
(Groupe d'Etude Sous-Marine de l'Atlantique)






1.1 Sensors


GPS (Global positioning system), only at the surface.

$$\begin{aligned} t_0 &= 6000 \text{ s}, & \ell^0 &= (-4.4582279^\circ, 48.2129206^\circ) \pm 2.5m \\ t_f &= 12000 \text{ s}, & \ell^f &= (-4.4546607^\circ, 48.2191297^\circ) \pm 2.5m \end{aligned}$$

Sonar (KLEIN 5400 side scan sonar).










Screenshot of SonarPro




Mine detection with SonarPro

Loch-Doppler returns the speed robot $\mathbf{v}_r$.

$$\mathbf{v}_r \in \tilde{\mathbf{v}}_r + 0.004 * [-1, 1] . \tilde{\mathbf{v}}_r + 0.004 * [-1, 1]$$

Inertial central (Octans III from IXSEA).

$$\begin{pmatrix} \phi \\ \theta \\ \psi \end{pmatrix} \in \begin{pmatrix} \tilde{\phi} \\ \tilde{\theta} \\ \tilde{\psi} \end{pmatrix} + \begin{pmatrix} 1.75 \times 10^{-4} \cdot [-1, 1] \\ 1.75 \times 10^{-4} \cdot [-1, 1] \\ 5.27 \times 10^{-3} \cdot [-1, 1] \end{pmatrix}.$$



Six mines have been detected.

i	0	1	2	3	4	5
$\tau(i)$	7054	7092	7374	7748	9038	9688
$\sigma(i)$	1	2	1	0	1	5
$\tilde{r}(i)$	52.42	12.47	54.40	52.68	27.73	26.98

6	7	8	9	10	11
10024	10817	11172	11232	11279	11688
4	3	3	4	5	1
37.90	36.71	37.37	31.03	33.51	15.05

Exercise. Draw the association graph associated with the detections.

1.2 Constraints

$$t \in \{6000.0, 6000.1, 6000.2, \dots, 11999.4\},$$

$$i \in \{0, 1, \dots, 11\},$$

$$\begin{pmatrix} p_x(t) \\ p_y(t) \end{pmatrix} = 111120 \begin{pmatrix} 0 & 1 \\ \cos\left(\ell_y(t) * \frac{\pi}{180}\right) & 0 \end{pmatrix} \begin{pmatrix} \ell_x(t) - \ell_x^0 \\ \ell_y(t) - \ell_y^0 \end{pmatrix}$$

$$\mathbf{p}(t) = (p_x(t), p_y(t), p_z(t)),$$

$$\mathbf{R}_\psi(t)=\begin{pmatrix}\cos\psi(t)&-\sin\psi(t)&0\\ \sin\psi(t)&\cos\psi(t)&0\\ 0&0&1\end{pmatrix},$$

$$\mathbf{R}_\theta(t)=\begin{pmatrix}\cos\theta(t)&0&\sin\theta(t)\\ 0&1&0\\ -\sin\theta(t)&0&\cos\theta(t)\end{pmatrix},$$

$$\mathbf{R}_\varphi(t) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \varphi(t) & -\sin \varphi(t) \\ 0 & \sin \varphi(t) & \cos \varphi(t) \end{pmatrix},$$

$$\mathbf{R}(t) = \mathbf{R}_\psi(t)\mathbf{R}_\theta(t)\mathbf{R}_\varphi(t),$$


$$\dot{\mathbf{p}}(t) = \mathbf{R}(t).\mathbf{v}_r(t),$$


$$||\mathbf{m}(\sigma(i))-\mathbf{p}(\tau(i))||\;=\;r(i),$$


$$\mathbf{R}^\top(\tau(i))\left(\mathbf{m}(\sigma(i))-\mathbf{p}(\tau(i))\right)\in [0]\times [0,\infty]^{\times 2},$$


$$m_z(\sigma(i))-p_z(\tau(i))-a(\tau(i))\in [-0.5,0.5]$$

1.3 GESMI









Extended Kalman smoother

2 Intervals in lattices

2.1 Lattices

A *lattice* $(\mathcal{E}, \leq)$ is a partially ordered set, closed under least upper and greatest lower bounds.

The least upper bound of x and y is called the *join*: $x \vee y$.

The greatest lower bound is called the *meet*: $x \wedge y$.

The Cartesian product of two lattices $(\mathcal{E}_1, \leq_1)$ and $(\mathcal{E}_2, \leq_2)$ is a lattice $(\mathcal{E}, \leq)$ with

$$(a_1, a_2) \leq (b_1, b_2) \Leftrightarrow ((a_1 \leq_1 b_1) \text{ and } (a_2 \leq_2 b_2)).$$

Exercise. $\mathcal{L} = ((\mathbb{B}, \mathbb{R}), \leq)$ is a lattice.

$$(\text{false}, 5) \vee (\text{true}, 2) = ?$$

$$(\text{false}, 5) \wedge (\text{true}, 2) = ?$$

$$\top(\mathcal{L}) = ?$$

$$\perp(\mathcal{L}) = ?$$

Example. The set $(\mathbb{R}^n, \leq)$ is a lattice with

$$\mathbf{x} \leq \mathbf{y} \Leftrightarrow \forall i \in \{1, \dots, n\}, x_i \leq y_i.$$

Example.

The powerset $\mathcal{P}(\mathbb{E})$ of all subsets of $\mathbb{E}$ is a lattice with respect to the inclusion $\subset$.

What is the meet ? What is the join ?

Example

The set $\mathcal{F}$ of all functions from $\mathbb{R}$ to $\mathbb{R}^n$ is a lattice with

$$\mathbf{f} \leq \mathbf{g} \Leftrightarrow \forall t \in \mathbb{R}, \mathbf{f}(t) \leq \mathbf{g}(t)$$

An interval of $\mathcal{F}$ is called a *tube*.


Intervals. A *closed interval* (or *interval* for short) $[x]$ of a lattice $\mathcal{E}$ is a subset of $\mathcal{E}$ which satisfies

$$[x] = \{x \in \mathcal{E} \mid \wedge [x] \leq x \leq \vee [x]\} .$$

The set $\mathbb{I}\mathcal{L}$ of all intervals of a lattice $\mathcal{L}$ is also a lattice with respect to $\subset$.


Exercise. Draw the Hasse diagram of the set of Boolean interval $\mathbb{I}\mathbb{B}$.

Graph intervals





Both $\emptyset$ and $\mathcal{E}$ are intervals of $\mathcal{E}$.

Exercise. Contract the graph interval with respect to the constraint " $\mathcal{G}$ is an equivalence relation".




Interval in the lattice $(\mathcal{P}(\mathbb{R}^n), \subset)$





An interval function (or tube) and a set interval


2.2 Interval arithmetic







$[A], [B], [A] \cap [B], [A] \cup [B],$
 $[A] \setminus [B], ([A] \cup [B]) \setminus ([A] \cap [B]).$

Intersection.

$$\begin{aligned} [A] \sqcap [B] &= \{X, X \in [A] \text{ and } X \in [B]\} \\ &= [A^- \cup B^-, A^+ \cap B^+]. \end{aligned}$$





2.3 Contractors in lattices

A CSP is composed of a set of variables $\{x_1, \dots, x_n\}$, of constraints $\{c_1, \dots, c_m\}$ and of domains $\{\mathbb{X}_1, \dots, \mathbb{X}_n\}$.


The domains $\mathbb{X}_i$ should belong to a lattice $(\mathcal{L}_i, \subset)$.

For SLAM, the domains are

(i) intervals of $\mathbb{R}^n$ to represent the location of the marks,

(ii) tubes to represent the unknown trajectory and

(iii) intervals of subsets of $\mathbb{R}^n$ to represent the free space.



Example

$$\left\{ \begin{array}{l} \mathbb{A} \subset \mathbb{B} \\ \mathbb{A} \in [\mathbb{A}], \mathbb{B} \in [\mathbb{B}]. \end{array} \right.$$

Since

$$\mathbb{A} \subset \mathbb{B} \Leftrightarrow \mathbb{A} = \mathbb{A} \cap \mathbb{B} \Leftrightarrow \mathbb{B} = \mathbb{A} \cup \mathbb{B}.$$

the optimal contractor is

$$\left\{ \begin{array}{l} \text{(i)} \quad [\mathbb{A}] := [\mathbb{A}] \sqcap ([\mathbb{A}] \cap [\mathbb{B}]) \\ \text{(ii)} \quad [\mathbb{B}] := [\mathbb{B}] \sqcap ([\mathbb{A}] \cup [\mathbb{B}]) \end{array} \right.$$

Tarski theorem.

If $(\mathcal{L}, \leq)$ is a lattice and $f : \mathcal{L} \rightarrow \mathcal{L}$ is monotonic (i.e., $a \leq b \Rightarrow f(a) \leq f(b)$), then $x_{k+1} = f(x_k)$, converges to the greatest x_∞ such that

$$\begin{cases} x_\infty = f(x_\infty) & \text{(fixed point)} \\ x_\infty \leq x_0 \end{cases}$$


2.4 Propagation

Consider the following CSP


$$\left\{ \begin{array}{ll} \text{(i)} & \mathbb{X} \subset \mathbb{A} \\ \text{(ii)} & \mathbb{B} \subset \mathbb{X} \\ \text{(iii)} & \mathbb{X} \cap \mathbb{C} = \emptyset \\ \text{(iv)} & f(\mathbb{X}) = \mathbb{X}, \end{array} \right.$$

where $\mathbb{X} \subset \mathbb{R}^2$, f is a rotation of $-\frac{\pi}{6}$, and


$$\left\{ \begin{array}{ll} \mathbb{A} & = \left\{ (x_1, x_2), x_1^2 + x_2^2 \leq 3 \right\} \\ \mathbb{B} & = \left\{ (x_1, x_2), (x_1 - 0.5)^2 + x_2^2 \leq 0.3 \right\} \\ \mathbb{C} & = \left\{ (x_1, x_2), (x_1 - 1)^2 + (x_2 - 1)^2 \leq 0.15 \right\} \end{array} \right.$$




(a)




(b)




(c)




(d)




(e)



(f)



(g)



(h)

$$(a) \quad [A]$$

$$(b) \quad [B]$$

$$(c) \quad [C]$$

$$(d) \quad X \subset A$$

$$(e) \quad B \subset X$$

$$(f) \quad X \cap C = \emptyset$$

$$(g) \quad f(X) = X$$


$$(h) \quad (f(X) = X)^\infty$$

3 Range-only SLAM with occupancy maps


$$\begin{cases} \dot{\mathbf{x}}(t) = \mathbf{f}(\mathbf{x}(t), \mathbf{u}(t)) & \text{(evolution equation)} \\ z(t) = d(\mathbf{x}(t), \mathbb{M}) & \text{(map equation)} \end{cases}$$

where $t \in \mathbb{R}$, $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{u} \in \mathbb{R}^m$, $\mathbb{M} \in \mathcal{C}(\mathbb{R}^q)$ is the occupancy map.

Unknown: the map $\mathbb{M}$ and the trajectory $\mathbf{x}$.




Impact, covering and dug zones





Tescase


$$\begin{cases} \dot{x}_1(t) &= u_1(t) \cos(u_2(t)) \\ \dot{x}_2(t) &= u_1(t) \sin(u_2(t)) \\ z(t) &= d(\mathbf{x}(t), \mathbb{M}). \end{cases}$$



Actual trajectory and dug space








Width of the tubes $[\mathbf{x}](t)$

4 Range only SLAM with undistinguishable marks

$$\begin{cases} \dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{u}) & \text{(evolution equation)} \\ (t_i, \mathcal{H}_i(\mathbf{x})) & \text{(sector list)} \end{cases}$$

Example. A robot is located at (x_1, x_2) . If at time t_3 the robot detects one single mark at a distance $d \in [4, 5]\text{m}$,


$$\mathcal{H}_3(\mathbf{x}) : \left\{ \mathbf{a} \in \mathbb{R}^2 \mid (x_1 - a_1)^2 + (x_2 - a_2)^2 \in [16, 25] \right\}.$$



The robot has detected the mark inside the ring


Theorem. Consider a set of marks $\mathcal{M} \subset \mathbb{R}^q$. Define the free space as $\mathbb{F} = \{\mathbf{p} \in \mathbb{R}^q \mid \mathbf{p} \notin \mathcal{M}\}$. Consider m sectors $\mathbb{H}_1, \dots, \mathbb{H}_m$, each of them containing exactly one mark and define $\mathbf{a}(i) = \mathcal{M} \cap \mathbb{H}_i$. We have

- (i) $\mathbb{H}_i \subset \mathbb{H}_j \Rightarrow \mathbf{a}(i) = \mathbf{a}(j)$
- (ii) $\mathbb{H}_i \cap \mathbb{H}_j = \emptyset \Rightarrow \mathbf{a}(i) \neq \mathbf{a}(j)$
- (iii) $\mathbb{H}_i \subset \mathbb{H}_j \Rightarrow \mathbb{H}_j \setminus \mathbb{H}_i \subset \mathbb{F}$.




Each of the two black zones contains a single mark and that no mark exists in the hatched area.


- (i) $\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{u})$
- (ii) $\mathbb{H}_i = \mathcal{H}_i(\mathbf{x}(t_i))$
- (iii) $\mathbf{a}(i) \in \mathbb{H}_i$
- (iv) $\mathbf{a}(i) = \mathbf{a}(j) \Leftrightarrow g_{ij} = \mathbf{1}$
- (v) $\mathbf{a}(i) \in \mathbb{H}_j \Leftrightarrow g_{ij} = \mathbf{1}$
- (vi) $g_{ij} = \mathbf{1} \Rightarrow \mathbb{H}_j \setminus \mathbb{H}_i \subset \mathbb{F}$
- (vii) $\mathbf{a}(i) \notin \mathbb{F}$




Contractor graph

4.1 Testcase







Free space $\mathbb{F}$.